

# Universal Methylated Mouse DNA Standard & Control Primers

Cat. Nos. D5012

Storage: -20 °C



ZYMO RESEARCH

The Beauty of Science is to Make Things Simple


## Product Information

### Product Contents:

	Cat. # D5012	Storage Temp.
Universal Methylated Mouse DNA Standard	5 µg/20 µl	-20 °C
mMLH1 Primers	20 µl	-20 °C

### Description:

The **Universal Methylated Mouse DNA Standard** includes enzymatically methylated DNA together with a specially-designed primer set to be used in conjunction with Zymo Research Corporation's **EZ DNA Methylation™**, **EZ DNA Methylation-Gold™**, and **EZ DNA Methylation-Direct™** kits to assess the efficiency of bisulfite-mediated conversion of DNA. The supplied DNA was isolated from male mouse strain Balb/c, and is enzymatically methylated at all cytosine positions comprising CG dinucleotides by M.Sss1 methyltransferase<sup>1</sup> (EC 2.1.1.37; Figure 1).



**Figure 1.** M.Sss1 methyltransferase methylates all cytosine residues in the double-stranded CpG context.

The primer set is designed to amplify a fragment of the mouse MLH1 mismatch repair gene following bisulfite treatment. The methylated cytosines comprising CG dinucleotides remain unconverted following bisulfite treatment, whereas non-methylated cytosines are converted into uracil and detected as thymine after PCR.

### References:

1. Nur *et al.* J. Bacteriol. 164: 19-24 (1985).

### Protocol:

*Note: We recommend using ZymoTaq™ DNA Polymerase or other hot-start DNA polymerases for amplification of bisulfite-treated DNA.*

#### 1. PCR Setup:

The following setup is designed for a 25 µl total reaction volume:

Component	Volume	Final Conc.
mMLH1 primer I*	Variable	0.2 to 0.8 µM
mMLH1 primer II*	Variable	0.2 to 0.8 µM
Bisulfite-converted DNA**	2 µl	up to 20 ng/µl
10 mM dNTP mix	0.5 µl	0.2 mM each dNTP
Standard PCR buffer	Variable	1x
MgCl <sub>2</sub> or MgSO <sub>4</sub>	Variable	1-4 mM, if needed
ZymoTaq™ DNA Polymerase (or other Hot-start DNA polymerase)	Variable	1 to 2 units
<b>Add water to 25 µl</b>		

\* Alternatively, you may substitute primers of your choice.

\*\* Remember to bisulfite-treat the DNA prior to performing PCR.

#### 2. Recommended Thermocycler Conditions:

- 95 °C, 10 minutes
- 95 °C, 30 seconds
- 58 °C, 30 to 60 seconds
- 72 °C, 60 seconds
- Repeat steps B through D an additional 29 to 39 times depending on the polymerase used.
- 72 °C, 7 minutes
- 4 °C

The PCR amplicon can now be used directly for sequencing analysis or cloning.

### Product Specifications:

- Universal Methylated DNA Standard, 5 µg/20 µl.  
Source: DNA isolated from male mouse strain Balb/c [enzymatically methylated by M.Sss1 Methyltransferase (EC 2.1.1.37)].  
Concentration: 250 ng/µl in TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0).  
Storage: -20 °C
- Control Primers.  
Concentration: 20 µM each primer in TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0)  
Volume: 20 µl of mixed primers  
Storage: -20 °C  
Sequence:

mMLH1 Primer I:

5' - GGTGTACGAAGTTATTTTTATTTTAGTC - 3'

mMLH1 Primer II:

5' - ACCCAACGATACCTAATAATAAAACC - 3'

### Appendix:

The expected PCR amplicon for the Universal Methylated Mouse DNA Standard is 304 bp, corresponding to nucleotide positions 430 to 778 of mouse MLH1 DNA including the regions (italicized) that hybridize to the primers (GenBank Accession #: AF400617).

Original sequence of mouse MLH1 DNA for bisulfite treatment and PCR amplification (sense strand 5' to 3'). The cytosines (underlined) in the CpG dinucleotide context (bold capitol letters) are methylated enzymatically by M.Sss1 methyltransferase:

```

421 -----g gtgtaCGaag tcaccctcac cccagcCGCG
461 acccttcaag gccaaagaagC GgcagaggC Gaggcctgcc
501 CGCGtCGctc tctcctcCGg agtgagcaCG gCGgccaaaag
541 acatgtcacc ctgcCGcaga CGctCGacca gggcCGCGCG
581 ttctctCGtcc cctacaaacC GctCGtagaa ttCGtgctCG
621 gcctCGtagt ggCGcctcaC GtCGCGttcc CGagtagagg
661 CGaccaggCG gCGacacacc aggcacaggg cccCGtacc
701 CtcCGcaggc tccaccacca ggtatCGctg ggt-----
  
```

Continued on next page...

ZYMO RESEARCH CORP.

Phone: (949) 679-1190 ▪ Toll Free: (888) 882-9682 ▪ Fax: (949) 266-9452 ▪ info@zymoresearch.com ▪ www.zymoresearch.com

**Appendix (continued...):**

Expected sequence of the above DNA following bisulfite treatment.  
Methylated cytosines in the CpG dinucleotide context remain unconverted following bisulfite treatment, whereas non-methylated cytosines, or cytosines not in the CpG context, are converted to uracil and detected as thymine after PCR.

```

421  -----g gtgtaCGaag ttatTTTTat ttagtCGCG
461  Atttttaaag gtaagaagC GtagaggtC Gaggtttgtt
501  CGCGtCGttt ttttttCGg agtgagtaCG gCGgttaaag
541  atatgttatt ttgtCGtaga CGttCGatta gggTCGCGCG
581  tttttCGttt ttataaaatC GttCGtagaa ttCGtgttCG
621  gtttCGtagt ggCGttttaC GtCGCGttt CGagtagagg
661  CGattaggCG gCGatatatt aggtataggg tttCGttatt
701  tttCGtaggt tttattatta ggtatCGtg ggt-----
    
```

**Trademarks and Disclaimers:**

™ Trademarks of Zymo Research Corporation.

This product is for research use only and should only be used by trained professionals. Wear protective gloves and eye protection. Follow the safety guidelines and rules enacted by your research institution or facility.

The Polymerase Chain Reaction (PCR) process is covered by U.S. Patent: #4,683,195; 4,683,202 assigned to Hoffmann-La Roche. Patents pending in other countries. No license under these patents to use the PCR process is conveyed expressly or by implication to the purchaser by the purchase of Zymo Research's products. Further information on purchasing licenses to practice the PCR process can be obtained from the director of Licensing at Applied Biosystems, 850 Lincoln Centre Drive, Foster City, California 94404 or at Roche Molecular Systems, Inc., 1145 Atlantic Avenue, Alameda, California 94501.

Version 2.1.6

**Also Available:**

Product Name	Size	Catalog number
EZ DNA Methylation™ Kit	50	D5001
	200	D5002
	2 x 96	D5003
	2 x 96	D5004
EZ DNA Methylation-Gold™ Kit	50	D5005
	200	D5006
	2 x 96	D5007
	2 x 96	D5008
EZ DNA Methylation-Direct™ Kit	50	D5020
	200	D5021
	2 x 96	D5022
	2 x 96	D5023
EZ DNA Methylation-Startup™ Kit	1 Kit	D5024
EZ Bisulfite DNA Clean-up Kit™	50	D5025
	200	D5026
	2 x 96	D5027
	2 x 96	D5028
Universal Methylated DNA Standard	1 set	D5010
Universal Methylated Human DNA Standard	1 set	D5011
Human HCT116 DKO Methylation Standards	1 set	D5014
Human HCT116 DKO Non-methylated DNA Standard	5 µg	D5014-1
Human HCT116 DKO Methylated DNA Standard	5 µg	D5014-2
Bisulfite Converted Universal Methylated Human DNA Standard	1 set	D5015
<i>E. coli</i> Non-methylated Genomic DNA	5 µg	D5016
ChIP DNA Clean & Concentrator™	50	D5201
	50	D5205
Methylated-DNA IP Kit	10	D5101
Anti-5-Methylcytosine Monoclonal Antibody (clone 10G4)	50 µg	A3001-50
	200 µg	A3001-200
Zymo <i>Taq</i> ™ DNA Polymerase	50	E2001
	200	E2002
Zymo <i>Taq</i> ™ PreMix (2X concentrated)	50	E2003
	200	E2004
CpG Methylase (M.SssI)	200 units	E2010
	400 units	E2011

**ZYMO RESEARCH CORP.**